



Xenin and Related Peptides: Potential Therapeutic Role in Diabetes and Related Metabolic Disorders

Craig, S. L., Irwin, N., & Gault, V. A. (2021). Xenin and Related Peptides: Potential Therapeutic Role in Diabetes and Related Metabolic Disorders. *Clinical Medicine Insights: Endocrinology and Diabetes*, 14, 117955142110438. [11795514211043868]. <https://doi.org/10.1177/11795514211043868>

[Link to publication record in Ulster University Research Portal](#)

Published in:

Clinical Medicine Insights: Endocrinology and Diabetes

Publication Status:

Published (in print/issue): 01/01/2021

DOI:

[10.1177/11795514211043868](https://doi.org/10.1177/11795514211043868)

Document Version

Publisher's PDF, also known as Version of record

General rights

Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact pure-support@ulster.ac.uk.

Xenin and Related Peptides: Potential Therapeutic Role in Diabetes and Related Metabolic Disorders

Sarah L Craig , Nigel Irwin and Victor A Gault

Faculty of Life and Health Sciences, School of Biomedical Sciences, Ulster University, UK.

Clinical Medicine Insights:
Endocrinology and Diabetes
Volume 14: 1–10
© The Author(s) 2021
DOI: 10.1177/11795514211043868



ABSTRACT: Xenin bioactivity and its role in normal physiology has been investigated by several research groups since its discovery in 1992. The 25 amino acid peptide hormone is secreted from the same enteroendocrine K-cells as the incretin hormone glucose-dependent insulinotropic polypeptide (GIP), with early studies highlighting the biological significance of xenin in the gastrointestinal tract, along with effects on satiety. Recently there has been more focus directed towards the role of xenin in insulin secretion and potential for diabetes therapies, especially through its ability to potentiate the insulinotropic actions of GIP as well as utilisation in dual/triple acting gut hormone therapeutic approaches. Currently, there is a lack of clinically approved therapies aimed at restoring GIP bioactivity in type 2 diabetes mellitus, thus xenin could hold real promise as a diabetes therapy. The biological actions of xenin, including its ability to augment insulin secretion, induce satiety effects, as well as restoring GIP sensitivity, earmark this peptide as an attractive antidiabetic candidate. This minireview will focus on the multiple biological actions of xenin, together with its proposed mechanism of action and potential benefits for the treatment of metabolic diseases such as diabetes.

KEYWORDS: Xenin-25, glucose-dependent insulinotropic polypeptide, insulin secretion, satiety, hybrid peptides, diabetes

RECEIVED: March 24, 2021. **ACCEPTED:** August 9, 2021.

TYPE: Review Article

FUNDING: The author(s) received no financial support for the research, authorship, and/or publication of this article.

DECLARATION OF CONFLICTING INTERESTS: The author(s) declared the following potential conflicts of interest with respect to the research, authorship, and/or publication of

this article: The Editor in Chief of Clinical Medicine Insights: Endocrinology and Diabetes is an author of this paper, therefore, the peer review process was managed by alternative members of the Board and the submitting Editor had no involvement in the decision-making process.

CORRESPONDING AUTHOR: Nigel Irwin, Faculty of Life and Health Sciences, School of Pharmacy and Pharmaceutical Sciences, Ulster University, Coleraine Campus, Cromore Road, Coleraine BT52 1SA, Northern Ireland. Email: n.irwin@ulster.ac.uk

Introduction

Originally identified from human gastric duodenal and jejunal mucosal isolates,¹ xenin, a naturally occurring 25-amino acid peptide, is synthesised from its 35-amino acid (aa) precursor pro-xenin.^{2–4} Interestingly, all 35-aa residues of yeast and mammalian alpha coat protein (COPA) are identical to that of pro-xenin.³ Biologically active xenin-25 (otherwise termed xenin) is then released following the action of pepsin on pro-xenin.^{5,6} Xenin has long been recognised as the human equivalent of the amphibian peptide xenopsin.⁷ Subsequent studies following on from original work by Feurle et al¹ that evidenced xenin in human gastric mucosa, demonstrate that xenin can be further extracted from the gut of various other species including dog, rabbit, rat and pig.^{6,8} In keeping with the view that the gut harbours numerous important regulatory peptide hormones, the highest concentrations of xenin are found within the gastrointestinal system.⁸ In this regard, xenin is synthesised and secreted into the circulation from a subpopulation of chromogranin A-positive enteroendocrine K-cells,⁹ along with the incretin hormone, GIP, in response to food ingestion. However, Hamscher et al⁸ also identified xenin in other key organs in dogs, including hypothalamus, liver, kidney, heart, pancreas, testes and skin. More recent studies have also identified xenin immunoreactivity within the endocrine pancreas,¹⁰ suggesting local production and biological activity in this organ.

Function, Potential Mechanism of Action and Therapeutic Application of Xenin

Xenin possesses numerous important biological actions that have been established in various animal models, (see Figure 1;

Table 1) which have previously been reviewed in depth.^{4,6} Briefly, key biological actions of xenin include control of energy balance and gastric transit,^{1,6,11,12} delay of gastric emptying in humans,¹³ appetite suppression,^{6,13–16} as well as regulating pancreatic exocrine and endocrine function.^{1,4,6,9,16–22} Xenin has also been shown to play a role in regulating normal bone physiology, potentially through indirect neural effects.²³ Studies have also clearly revealed that xenin can potentiate the insulin-releasing capabilities of GIP (Figure 2), the incretin hormone co-secreted with xenin from intestinal K-cells,^{19,21,24–26} highlighting favourable attributes for the treatment of diabetes. Despite this established biological profile, a specific xenin receptor has yet to be identified. There is a suggestion that aspects of the biological actions of xenin may be mediated through activation of the neurotensin receptor, due to structural similarities between the 2 peptides.²⁷ However, effects of xenin independent of neurotensin receptor activation have been demonstrated,²⁸ highlighting the need for further detailed studies in this area. Finally, although there is no direct evidence for xenin induced benefits in type 1 diabetes mellitus, reduction of beta-cell apoptosis¹⁰ alongside positive actions on islet cell transdifferentiation,²⁹ could be suggestive of positive effects of xenin in this disease state.

GIP potentiation

Resistance to the biological actions of GIP is a hallmark of type 2 diabetes mellitus, with the GIP-mediated incretin effect being severely diminished in people with diabetes (Figure 2).³⁴ However, despite the well-known importance of the incretin



Creative Commons CC BY: This article is distributed under the terms of the Creative Commons Attribution 4.0 License (<https://creativecommons.org/licenses/by/4.0/>) which permits any use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).




Figure 1. Representation of the main biological actions of xenin. The impact of xenin on adipose tissue, brain, pancreas and gastrointestinal tract are considered.

Table 1. Summary of evidence to support the main biological actions of xenin represented in Figure 1.

SPECIES	TREATMENT	MAIN OUTCOMES	REFERENCES
Gastrointestinal actions			
Rodent	Experimental design <ul style="list-style-type: none"> Dunkin-hartley guinea pigs Maximal efficacy of xenin-25 – 10^{-6} M 	<ul style="list-style-type: none"> In the jejunum <ul style="list-style-type: none"> Small relaxation followed by a large contraction In the colon <ul style="list-style-type: none"> Myokinetic relaxation effect 	Feurle et al ³⁰
Rodent	Experimental design <ul style="list-style-type: none"> Xenin-25 (1 μM) at 15-20 min intervals 	<ul style="list-style-type: none"> Relaxation of rat ileum 	Clemens et al ²⁷
Human	Experimental design <ul style="list-style-type: none"> Constant intravenous infusions: 0-300 min Infusion rates <ul style="list-style-type: none"> Xenin @ 4 pmol/kg infusion Xenin @ 12 pmol/kg – administered at the same relative flow rates as above 	<ul style="list-style-type: none"> Delay of gastric emptying in humans with and without T2DM Reduction in postprandial glucose levels 	Chowdhury et al ¹³
Anorexigenic effects			
Chick	Experimental design <ul style="list-style-type: none"> Central effects on feeding: <ul style="list-style-type: none"> Intracerebroventricular (ICV) injection of 0.75, 1.5 or 3.0 μg xenin. Peripheral effects on feeding: <ul style="list-style-type: none"> Intraperitoneal injection of avian saline, 0.2, 2.0 or 20.0 μg xenin dissolved in in 180 min fasted chicks Gastrointestinal transit rate: <ul style="list-style-type: none"> Non-fasted chicks received the same ICV treatments as above. Immediately after injection, chick was gavaged with feed slurry at a mass of 4.0% body weight 	<ul style="list-style-type: none"> Anorexigenic actions and delay gastrointestinal transit rate in chicks 	Cline et al ¹¹

(Continued)

Table 1. (Continued)

SPECIES	TREATMENT	MAIN OUTCOMES	REFERENCES
Rodent	Experimental design <ul style="list-style-type: none">Fasted (16 h) mice re-fed with pre-weighed food pellet for 1 hMice then given intraperitoneal injection of saline, xenin (50 µg/g bw) or urocortin (3 nmol/mouse)Rate of gastric emptying was calculated as follows: Gastric emptying (%) = {1 – (wet weight of food recovered from the stomach/wet weight of food intake)} × 100¹²The effect of xenin on food intake was examined in ad libitum-fed wild-type mice. Mice were injected intraperitoneally with xenin (50 µg/g bw) or saline and cumulative food intake was measured 1, 2, 4, 6, 8, 12, 18 and 24 h after injection¹⁵Mice were fasted for 12 h before subcutaneous injection of 50, 100 or 500 nmol/kg xenin. Mice were then allowed free access to normal chow. Cumulative food intake was measured at 30, 60, 60 and 120 min post injection¹⁸	<ul style="list-style-type: none">Reduction of gastric emptying by 93% and induction of satiety	Kim and Mizuno, ¹² Alexiou et al, ¹⁴ Leckstrom et al, ¹⁵ Taylor et al, ¹⁸ Cooke et al, ³¹ and Bhavya et al ³²
Adipose Tissue			
Rodent	Experimental design <ul style="list-style-type: none"><i>Ad libitum</i> fed mice received 2 ICV injections of xenin (5 µg) at 10:00 h and 22:00 hBody weight and food weight were measured immediately prior to the first injection and 24 h after the first injection. Mice were euthanised 12 to 14 h after the second injection Epididymal adipose tissues and skeletal muscles were collected for RNA and protein analyses	<ul style="list-style-type: none">Increased expression of lipolytic markers	Bhavya et al ³²
3T3-L1 mouse adipocyte cell line	Experimental design <ul style="list-style-type: none">Immortalised 3T3-L1 fibroblasts differentiated 2 days post confluence in the absence or presence of xenin-25-Gln (10⁻⁶ M). Test peptides were added only during the key growth phase when the differentiation cocktail was presentGlycerol release, glucose uptake and gene expression were assessed	<ul style="list-style-type: none">Increased glycerol releaseKey adipogenic and lipolytic genes upregulatedStimulated insulin-induced glucose uptake	English et al ³³




Figure 2. Representation of the incretin effect mediated by GLP-1 and GIP under normal and diabetic conditions, with perceived xenin benefits in diabetes. (a) The incretin response under normal physiology alongside (b) the perturbed incretin response in T2DM, with (c) xenin acting as a GIP potentiator to restore GIP sensitivity in T2DM.

effect to regulate normal blood glucose levels,³⁵ established treatments for type 2 diabetes fail to address this issue. Indeed, incretin-based therapeutics focus largely on augmenting the biological actions of the sister incretin glucagon-like peptide-1 (GLP-1). However, recent exciting clinical findings with a

dual-acting GLP-1 and GIP receptor hybrid peptide exhibiting strong bias towards the GIP receptor,³⁶ suggests that GIP resistance in type 2 diabetes is surmountable. In this regard, xenin has been shown to potentiate the insulinotropic actions of GIP in rodent models of diabetes.^{18-22,24-26} Whilst the

precise mechanism of xenin-induced GIP potentiation remains to be fully elucidated,^{25,27,37} it may be linked to acetylcholine M3 receptor signalling on pancreatic beta cells.²⁵ However, there is also good evidence for a direct effect of xenin on beta cells,⁶ that is reinforced by knowledge that xenin is produced and secreted locally within islets.¹⁰

Appetite suppression

Several studies have demonstrated the role of xenin in regulating energy intake. Administration of xenin reduces calorie consumption and delays gastric emptying in mice, rats, chicks and humans^{6,11,13–15,18,31} suggesting xenin may act directly on the gastrointestinal tract to induce satiety. This effect may occur through receptor binding at nerve terminal ends, which then influences the nucleus of the solitary tract anorexigenic activity, or hypothalamic receptors involved in energy homeostasis.³⁸ Indeed, hypothalamic neurons appear to have direct involvement in regulation of calorie intake following intraperitoneal administration of xenin, suggesting centrally mediated effects.¹⁵ Interestingly, more recent studies have characterised xenin activity in both peripheral and central regions linked to regulating feeding in goldfish, to induce anorexigenic actions.³⁹ It has also been demonstrated that xenin, when administered intracerebroventricularly in rats or peripherally in mice, may act through CRH-dependent signalling pathways to regulate food intake.³⁸ However, it has been established that anorexic effects of xenin are independent of both the leptin- and melanocortin-dependent signalling pathways.¹⁵

Lipid metabolism

In addition to its role in reducing food intake, xenin has also been shown to cause alterations in the expression of genes involved in lipid metabolism, as well as proteins found within white adipose tissue.³² There was an original hypothesis that xenin acts on adipose tissue to stimulate lipolysis, and that xenin may hold promise as an anti-obesity therapy by reducing adipose fat depots, but such observations were somewhat inconsistent.³² Thus, English et al³³ recently revealed direct lipogenic and lipolytic actions of xenin in 3T3-L1 adipocytes, whilst also promoting adipocyte differentiation in 3T3-L1 pre-adipocytes, through alterations in gene expression of LPL and FASN, key promoters of 3T3-L1 differentiation.³³ The effects of xenin to positively modulate lipolysis, lipogenesis and adipocyte differentiation are likely modulated through NTR1 activation on the AKT/PI3K pathway.³³ However, it should be noted that the actions of xenin on lipid metabolism are still not well defined and require more detailed study, especially in light of some conflicting observations.^{32,33}

Pancreas

Immunoreactivity of xenin has been identified in human pancreatic extracts,⁸ where concentrations increased following

pepsin digestion.⁸ More recently, immunohistochemical-based methods demonstrated expression of xenin in both alpha- and beta cells, with both arginine and glucose acting as a stimulus for xenin secretion from the islet.¹⁰ Numerous biological roles of xenin in the pancreas have already been recognised, including secretion of insulin and glucagon, as well as effects of secretory activity in the exocrine pancreas.⁶ In addition, xenin exerts beneficial effects on beta cell growth and protection against apoptosis,⁶ with obvious therapeutic benefit in the context of diabetes. Moreover, recent studies in insulin-deficient *Ins1^{Cre/+}*; *Rosa26-eYFP* transgenic mice with islet cell lineage tracing capabilities reveal positive effects of xenin on islet cell differentiation, including maintenance of beta cell identity and prevention of beta cell de-differentiation.²⁹ These positive effects on islet cell architecture may be related to potentiation of the biological actions of GIP, since GIP has established benefits on beta cell growth and survival, as well as transdifferentiation.^{40–43} The mechanisms related to these xenin-mediated pancreatic islet actions are somewhat disputed however, with proposed importance of both direct and indirect actions. Thus, xenin has been shown to directly stimulate glucagon and insulin secretion *in vitro* when applied to cultured pancreatic alpha- and beta-cells, respectively.¹⁷ These direct receptor-mediated actions are strengthened by evidence of local xenin production and secretion within pancreatic islets.¹⁰ On the other hand, there are also reports to suggest that xenin does not directly enhance GIP-mediated insulin exocytosis, with these effects stimulated through activation of acetylcholine containing enteric neurons that are in direct contact with the pancreas.²⁵

Polycystic ovary syndrome

Insulin resistance is an established pathological feature of type 2 diabetes mellitus, with polycystic ovary syndrome (PCOS) also closely associated with obesity and insulin resistance.⁴⁴ Thus, similar to diabetes, previous research has defined a relationship between xenopsin-related-peptide-1 and PCOS, where the levels of xenopsin-related-peptide-1 were significantly elevated in PCOS patients when compared to controls.⁴⁵ In this regard, serum xenin concentrations are significantly elevated in women with PCOS compared to women with no menstrual cycle abnormalities.⁴⁶ However, as with diabetes,⁶ the precise impact of xenin in PCOS and its pathophysiology remains to be fully elucidated. When viewed together, the above diverse biological actions of xenin emphasise potential for targeting related pathways for the amelioration of insulin resistance and related disease such as diabetes and PCOS.

Truncated Xenin Peptides and Analogues

Naturally occurring peptides such as xenin have many therapeutic advantages over small molecules, including their diversity, safety, ease of synthesis, along with minimal risk of drug-drug interactions.⁴⁷ Naturally occurring peptides also have a high binding affinity towards a broad, but specific range of therapeutic targets and are often very potent, resulting in

Table 2. Amino acid sequences of xenin-25 as well as its related stable analogues and naturally occurring fragment peptides.

PEPTIDE	AMINO ACID SEQUENCE	REFERENCES
Xenin-25	M-L-T-K-F-E-T-K-S-A-R-V-K-G-L-S-F-H-P-K-R-P-W-I-L-OH	Feurle et al ¹
Xenin-25-Gln	M-L-T-Q-F-E-T-Q-S-A-Q-V-Q-G-L-S-F-H-P-Q-Q-P-W-I-L-OH	Parthsarathy et al ²²
Xenin-25[Lys ¹³ PAL]	M-L-T-K-F-E-T-K-S-A-R-V-K-(N-ε-(γ-GLU(hexadecanoyl))-G-L-S-F-H-P-K-R-P-W-I-L-OH	Gault et al ²¹
Xenin 9-25 (Xenin-17)	S-A-R-V-K-G-L-S-F-H-P-K-R-P-W-I-L-OH	Martin et al ²⁰
Xenin 11-25 (Xenin-15)	R-V-K-G-L-S-F-H-P-K-R-P-W-I-L-OH	Martin et al ²⁰
Xenin 14-25 (Xenin-12)	G-L-S-F-H-P-K-R-P-W-I-L-OH	Martin et al ²⁰
Xenin 18-25 (Xenin-8)	H-P-K-R-P-W-I-L-OH	Martin et al ²⁰
Xenin 18-25-Gln	H-P-Q-Q-P-W-I-L-OH	Martin et al ¹⁶
Xenin 20-25 (Xenin-6)	K-R-P-W-I-L-OH	Craig et al ²⁶ and Feurle et al ⁵⁰
Xenin-6-psi	K-(CH ₂ NH)-R-P-W-I-L-OH	Craig et al ²⁶ and Feurle et al ⁵⁰

enhanced efficacy, selectivity and specificity, even at lower therapeutic doses.⁴⁸ Therefore, peptide therapeutics are of great interest for drug developers. However, the clinical use of peptides is hindered by certain disadvantages, including their instability and susceptibility to enzymatic degradation, reduced oral bioavailability, limited cell membrane permeation and rapid renal clearance.⁴⁹ Fortunately, these limitations can be largely overcome through structural modification of the peptide,⁴⁹⁻⁵³ which has been demonstrated for xenin, as discussed below.

Stable analogues of xenin (Tables 2 and 3) with preserved or even enhanced bioactivity have been developed.⁶ Many of these xenin analogues possess notable beneficial metabolic effects in pre-clinical models of diabetes-obesity,^{16,22,54} which has been reviewed in detail previously.⁶ However, the use of truncated peptide fragments of xenin that retain the full biological actions of the parent peptide, could enhance therapeutic promise by making peptide synthesis easier and cheaper, as well as facilitating possible non-injectable peptide drug delivery.^{55,56} An earlier comprehensive exploration identified the degradation profile of xenin in mouse plasma, revealing the following C-terminally truncated metabolites; xenin 9-25, xenin 11-25, xenin 14-25 and xenin 18-25 (where xenin 18-25 represents xenin-8).²⁰ Subsequent characterisation revealed that only xenin-8 possessed biological activity equivalent to the parent peptide.²⁰ Indeed, this truncated octapeptide has long been recognised as a naturally occurring and biologically active derivative of xenin,^{17,57,58} that retains full insulinotropic actions.²⁰ Furthermore, amino acid substitution of the Lys and Arg residues for Gln in xenin-8, resulted in production of a fully enzymatically stable octapeptide that retained full gluco-regulatory and antidiabetic actions as full-length xenin.¹⁶ Subsequent recent research has now confirmed bioactivity of xenin-6 (xenin 20-25) at the level of the endocrine pancreas.^{26,50} Moreover, modification of xenin-6

through introduction of a reduced pseudopeptide bond between amino acid residues Lys-20 and Arg-21, to create xenin-6-psi, further increased bioactivity of this truncated peptide.^{26,50} Intriguingly, xenin-6-psi exerted potent metabolic actions in diabetic rodents and prominently augmented the biological actions of the incretin hormone GIP.²⁶ Thus, it appears that the 6 C-terminal residues of xenin are sufficient to facilitate receptor binding and activation of the full repertoire of xenin cell signalling pathways.

Dual and Triple Acting Therapeutic Approaches That Incorporate Xenin Elements

As noted above, truncated xenin peptides retain bioactivity and have promising antidiabetic actions.^{20,26,50} However, in such a multi-factorial disease as type 2 diabetes mellitus, monotherapy does not appear to adequately control glycaemia over the longer-term. Thus, multi-targeting unimolecular hybrid peptides, designed to simultaneously modulate multiple signalling pathways are now thought to offer superior therapeutic efficacy than single targeted compounds.⁶⁰ Indeed, data emerging from recent clinical studies with a dual-acting GLP-1/GIP compound, Tirzepatide (LY3298176), developed by Lilly, with strong bias towards the GIP receptor, fully support this notion.⁶¹ Data from phase 1 and 2 studies were extremely promising, with the compound now entering SURPASS phase 3 clinical trials to determine long-term efficacy and safety.⁶² Initial proof-of-concept for utilisation of multi-acting hybrid peptides comes from the naturally occurring dual agonist oxyntomodulin (OXM), that activates both GLP-1 and glucagon receptor pathways.⁶³ More recent studies demonstrate the opportunity of linking together individual bioactive peptide domains of different peptides, or engineering unique amino acid sequences that incorporate binding capabilities of 2 or more regulatory peptides, to create multi-targeting hybrid peptides.⁶⁴⁻⁶⁶

Table 3. Summary of study design and main experimental outcomes from studies with fragment peptides of xenin-25.

SPECIES	TREATMENT	MAIN OUTCOMES	REFERENCES
In vitro and rodent	<p>Experimental design</p> <ul style="list-style-type: none"> For food intake studies, fasted (18 h) mice were given intraperitoneal (i.p.) injections of either saline vehicle (0.9% w/v NaCl), xenin-8 or xenin-8-Gln at a dose of 500 nmol/kg bw. Cumulative food intake measured over 120 min For glucose homeostasis and insulin secretory studies, blood glucose and plasma insulin concentrations were measured immediately prior to and 15, 30 and 60 min after i.p. administration of glucose alone (18 mmol/kg bw) or in combination with either xenin 18 to 25 or xenin 18-25 Gln (each at 25 nmol/kg bw) in non-fasted mice 	<ul style="list-style-type: none"> Concentration-dependently stimulated insulin secretion Enhanced glucose-induced insulin release 	Martin et al ¹⁶
Rodent	<p>Experimental design</p> <ul style="list-style-type: none"> Twice daily i.p. injections of saline vehicle, xenin-8 or xenin-8-Gln (both at 25 nmol/kg bw) for 21 days in HFF mice Energy intake, body weight, non-fasting blood glucose and plasma insulin concentrations were assessed during the 21 days At the end of the treatment period, i.p. glucose tolerance (18 mmol/kg bw), biological response to GIP (18 mmol/kg glucose in combination with native GIP (25 nmol/kg); i.p.) and insulin sensitivity (15 U/kg bw; i.p.) tests were performed 	<ul style="list-style-type: none"> Both treatment regimens <ul style="list-style-type: none"> Elevated circulating plasma insulin concentrations Improved insulin sensitivity Xenin-8-Gln <ul style="list-style-type: none"> Improved glucose tolerance Augmented GIP-mediated glucose-lowering and insulin-releasing effects 	Martin et al ¹⁶
Rodent	<p>Experimental design</p> <ul style="list-style-type: none"> For food intake studies, fasted (18 h) lean mice were given intraperitoneal (i.p.) injections of either saline vehicle (0.9% w/v NaCl) or Ψ-xenin-6 at a dose of 25 or 250 nmol/kg bw. Cumulative food intake was measured at 30 min intervals for 180 min For acute effects of peptides on glucose tolerance and insulin secretion, blood glucose and plasma insulin concentrations were determined immediately prior to and 15, 30, 60 and 105 min after i.p. injection of glucose alone (18 mmol/kg bw) or in combination with test peptides (25 nmol/kg bw), as well as test peptides together with GIP (25 nmol/kg bw) in 4 h fasted mice To assess duration of peptide action, mice were administered saline vehicle or test peptides (25 nmol/kg bw) at 2, 4, 8 or 12 h prior to an i.p. glucose challenge (18 mmol/kg bw) and blood glucose measured 	<ul style="list-style-type: none"> Significantly reduced glucose levels Enhanced glucose-induced insulin release Enhanced the glucose-lowering action of GIP Exhibited satiety actions 	Craig et al ²⁶
Rodent	<p>Experimental design</p> <ul style="list-style-type: none"> Oral sitagliptin phosphate monohydrate once daily (50 mg/kg bw), intraperitoneal (i.p.) Ψ-xenin-6 twice daily (25 nmol/kg bw) or a combination of both compounds for 18 days in HFF mice Energy intake, body weight, non-fasting blood glucose and plasma insulin concentrations were assessed at regular intervals At the end of the treatment period, i.p. glucose tolerance (18 mmol/kg bw; 18 h-fasted mice), insulin sensitivity (25 U/kg bovine insulin; i.p.; 18 h-fasted mice) and pyruvate tolerance (2 g/kg sodium pyruvate; i.p.; 18 h-fasted mice) tests were performed HOMA-IR, fasting glucose (mmol/L) \times fasting insulin (mU/L)/22.5, was also calculated as a surrogate marker of insulin resistance Terminal analyses included extraction of pancreatic tissue for determination of pancreatic insulin content. In addition, liver tissue was processed for hepatic gene expression by qPCR after total RNA extraction 	<p>Ψ-xenin-6 alone</p> <ul style="list-style-type: none"> Reduced weight gain Reduced glucose levels as well as improved glucose tolerance and insulin sensitivity. <p>Positive effects on pancreatic islet architecture</p> <p>Ψ-xenin-6 and sitagliptin:</p> <ul style="list-style-type: none"> Prominent benefits on circulating glucose and insulin levels Improvements in attenuating gluconeogenesis Benefits on pancreatic islet architecture Improved insulin sensitivity 	Craig et al ⁵⁹

With regards to type 2 diabetes mellitus, Gault et al⁶⁷ initially indicated that a GLP-1 and GIP preparation, that combined long-acting acylated version of the parent peptides, displayed enhanced glucose-lowering and insulinotropic actions in animal models of diabetes. This being despite earlier observations that combined administration of individual enzymatically stable, but non-acylated GIP and GLP-1 mimetics was not associated with benefits beyond that of either peptide alone,⁶⁸⁻⁷⁰ however this could be related to differences in treatment regimens or animal models employed. Following on from this, a triple acting hybrid peptide comprising GLP-1, GIP

and glucagon was developed that offered some improvements in preclinical models of obesity-diabetes when compared to parent peptides.⁷¹ In addition, 2 separate CCK/GLP-1 fusion peptides have been characterised revealing notable benefits on appetite suppression, insulinotropic effects as well as beta cell function and morphology.^{64,72} Furthermore, numerous other dual- and triple-acting hybrid peptides have been developed that clearly advocate the therapeutic benefits of single peptide-based drugs capable of positivity modulating more than 1 receptor pathway for the treatment of diabetes.^{65,66,73-75} Tschöp et al⁷⁵ demonstrated that novel unimolecular combination

Table 4. Amino acid sequences of xenin incorporated multi-acting hybrid peptides.

PEPTIDE	AMINO ACID SEQUENCE	REFERENCES
(DAla ²)GIP/xenin-8-Gln	Y-[DA]-E-G-T-F-I-S-D-Y-S-I-A-M-H-P-Q-Q-P-W-I-L-OH	Hasib et al ⁵⁴ and Pathak et al ⁷⁴
Exendin-4/xenin-8-Gln	H-G-E-G-T-F-T-S-D-L-S-K-Q-M-E-E-A-V-R-L-F-I-E-W-L-K-N- AEEAc – AEEAc-H-P-Q-Q-P-W-I-L-OH	Craig et al ⁷⁶
Exendin-4/gastrin/xenin-8-Gln	H-G-E-G-T-F-T-S-D-L-S-K-Q-M-E-E-A-V-R-L-F-I-E-W-L-K-N- AEEAc – AEEAc-Y-G-W-L-D-F- AEEAc – AEEAc-H-P-Q-Q-P-W-I-L-OH	Hasib et al ⁸¹
Exendin-4(Lys ²⁷ PAL)/gastrin/xenin-8-Gln	H-G-E-G-T-F-T-S-D-L-S-K-Q-M-E-E-A-V-R-L-F-I-E-W-L-K(γ -Glu-palm)-N-AEEAc-AEEAc-Y-G-W-L-D-F-AEEAc-AEEAc-H-P-Q-Q-P-W-I-L-OH	Hasib et al ⁷⁸

therapies have superior efficacy, compared to current therapeutic options, thus having potential to reverse obesity and type 2 diabetes.

In terms of incorporating xenin into multi-acting hybrid peptides (Tables 4 and 5), this was first demonstrated in 2017 through a GIP/xenin entity, namely (DAla²)GIP/xenin-8-Gln.⁵⁴ Subsequent work with (DAla²)GIP/xenin-8-Gln has highlighted that twice-daily administration in high fat fed mice for 28 days significantly reduced food intake and body weight, with associated reductions in circulating glucose concentrations and HbA_{1c} levels, whilst improving glucose tolerance and insulin sensitivity.⁷⁶ Similar, but somewhat less striking antidiabetic effects were noted in *db/db* mice given (DAla²)GIP/xenin-8-Gln, demonstrating that the positive antidiabetic actions are transferable across diverse aetiologies of type 2 diabetes mellitus.⁷⁶ Remarkably, the same study also demonstrated long-acting positive metabolic effects of (DAla²)GIP/xenin-8-Gln following 14-day cessation of treatment.⁷⁶ This could suggest positive metabolic reprogramming induced by co-activation of GIP and xenin receptor pathways in keeping with positive effects on beta cell function and integrity,^{6,29,43} and represents a potential benefit for future antidiabetic therapy. Such observations are extremely important moving towards the clinical setting given the complex aetiology and progressive nature of type 2 diabetes mellitus in humans.⁷⁷ Subsequent investigations characterised a novel GLP-1/xenin hybrid peptide (exendin-4/xenin-8-Gln) that exhibited positive antidiabetic actions in high fat fed mice,⁷⁸ highlighting positive effects of combined modulation of GLP-1 and xenin related signalling pathways in diabetes. Hasib et al⁷⁸ also demonstrated the potential of combined modulation of GLP-1, gastrin and xenin signalling pathways,^{78,79} which was superior to the previously described dual-acting fusion peptide incorporating GLP-1 and gastrin only, namely ZP3022.⁸⁰

More recent work has explored the possibility of Ψ -xenin-6 to enhance the antidiabetic efficacy of the established dipeptidyl peptidase-4 (DPP-4) inhibitor drug sitagliptin.⁵⁹ Multiple metabolic advantages of combined Ψ -xenin-6 and sitagliptin therapy were observed, including benefits on body weight, circulating glucose and insulin along with additional enhancements to reduce gluconeogenesis and improve pancreatic islet

architecture.⁵⁹ Additional related studies have demonstrated how specifically elevating xenin concentrations through use of the methionine aminopeptidase inhibitor 2, TNP-470, can also augment the antidiabetic efficacy of sitagliptin.⁸² Moreover, as well as increasing xenin secretion, TNP-470 is a putative anti-obesity agent,⁸³⁻⁸⁵ highlighting obvious benefits of this treatment modality in obesity-driven forms of diabetes. Given xenin has confirmed GIP-potentiating actions, the combination of therapies that increase xenin bioactivity alongside established DPP-4 inhibitor drugs clearly warrants further consideration as a novel therapeutic option in the management of type 2 diabetes mellitus in humans.

Concluding Remarks

This minireview highlights the diverse biological actions of xenin, as well as the therapeutic potential for xenin and related truncated metabolites for diabetes and related disorders. Future studies are required to fully understand the signalling pathways and mechanisms involved in the insulinotropic, GIP-potentiating and anorexigenic actions of xenin, as well as the role of xenin signalling within benefits of associated hybrid peptides. Clarification of whether or not a specific xenin receptor exists is key in this paradigm. Nevertheless, xenin possesses a promising therapeutic repertoire that may result in the development of a safe, effective, long-acting and cost-effective therapy for obesity-diabetes.

Due to the multifactorial nature of type 2 diabetes mellitus, monotherapy is often not an effective treatment option. Thus, combination therapy or hybrid peptides have the potential to emerge as leading therapeutic approaches for this disease. Both approaches show promise with xenin-based therapies, demonstrating obvious advantages over monotherapy that is highly favourable moving towards the clinic.^{54,76,78,79} However, future studies are required to fully understand the mechanisms and pathways associated with satiety effects, insulinotropic and GIP-potentiating actions to gain a better understanding of the role of xenin and overall therapeutic potential of these hybrid peptides. Further to this, recent studies have highlighted the stability and metabolic benefits of Ψ -xenin-6 alone,²⁶ and in combination with established anti-diabetic therapies.⁵⁹ To date, hybrid peptides that contain a xenin

Table 5. Summary of study design and main experimental outcomes from studies with xenin incorporated multi-acting hybrid peptides.

SPECIES	TREATMENT	MAIN OUTCOMES	REFERENCES
Rodent	Experimental design <ul style="list-style-type: none"> Twice daily i.p. injections of saline vehicle, (DAla²)GIP/xenin-8-Gln (25 nmol/kg bw), exendin-4 (25 nmol/kg bw), or a combination of both peptides for 28 days in HFF mice, followed by 14 days cessation of treatment Energy intake, body weight, non-fasting blood glucose and plasma insulin concentrations were assessed at regular intervals At the end of the treatment period, i.p. glucose tolerance (18 mmol/kg bw; 18 h-fasted mice) and insulin sensitivity (25 U/kg bovine insulin; i.p.; non-fasted mice) tests were performed. Metabolic responses to acute re-administration of respective treatment regimens together with glucose was also examined On day 28 observations were continued in a sub-group (n=6) of mice following cessation of treatment regimens for a further 14 days, with assessment of the same parameters as documented above 	(DAla ²)GIP/xenin-8-Gln <ul style="list-style-type: none"> Reduction in food intake, body weight, circulating glucose and HbA_{1c} Improved glucose tolerance and insulin sensitivity Improved pancreatic architecture 	Craig et al ⁷⁶
Rodent	Experimental design <ul style="list-style-type: none"> Twice daily i.p. injections of saline vehicle, (DAla²)GIP/xenin-8-Gln (25 nmol/kg bw), exendin-4 (25 nmol/kg bw), or a combination of both peptides for 28 days in <i>db/db</i> mice Energy intake, body weight, non-fasting blood glucose and plasma insulin concentrations were assessed at regular intervals At the end of the treatment period, i.p. glucose tolerance (18 mmol/kg bw; 18 h-fasted mice) and insulin sensitivity (50 U/kg bovine insulin; i.p.; non-fasted mice) tests were performed 	(DAla ²)GIP/xenin-8-Gln in combination with exendin-4 was required to induce beneficial effects on glucose tolerance, insulin sensitivity	Craig et al ⁷⁶
Rodent	Experimental design: <ul style="list-style-type: none"> Twice daily i.p. injections of saline vehicle (0.9% w/v NaCl), exendin-4, exendin-4/gastrin/xenin-8-Gln alone and in combination with (DAla²) GIP (each peptide at 25 nmol/kg bw) for 21 days in HFF mice Cumulative food intake, body weight, non-fasting glucose and insulin concentrations were monitored at regular intervals Circulating glucagon, amylase activity and blood lipid profile were assessed at the end of the treatment period Glucose tolerance (18 mmol/kg bw; i.p.), metabolic response to GIP (18 mmol/kg glucose in combination with native GIP (25 nmol/kg); i.p.) and insulin sensitivity (25 U/kg bw; i.p.) tests were performed at the end of the treatment period On day 21 locomotor activity and energy expenditure were assessed 	<ul style="list-style-type: none"> Reduced circulating glucose and increased plasma insulin concentrations Improved glucose tolerance, insulin sensitivity and metabolic response to GIP Reduced LDL-cholesterol and body fat mass Normalised pancreatic islet and beta-cell area Increase in energy expenditure and locomotor activity in mice treated with exendin-4/gastrin/xenin-8-Gln in combination with (DAla²)GIP 	Hasib et al ⁸¹
Rodent	Experimental design <ul style="list-style-type: none"> Dosing Regimen: Twice-daily injections of saline vehicle, exendin-4/gastrin, exendin-4/gastrin/xenin-8-Gln, or exendin-4(Lys²⁷PAL)/gastrin/xenin-8-Gln (each at 25 nmol/kg bw; ip) for 31 days in <i>ob/ob</i> mice Energy intake, body weight, non-fasting blood glucose and plasma insulin concentrations were assessed at regular intervals Plasma glucagon, amylase activity, 24-h glucose profile and whole blood HbA_{1c} were measured on day 31 At the end of the treatment period, glucose tolerance (18 mmol/kg bw; ip), metabolic response to GIP (18 mmol/kg glucose in combination with native GIP [25 nmol/kg]; ip) and insulin sensitivity (50 U/kg bw; ip) tests were conducted Percentage body fat and pancreatic insulin content were also determined 	<ul style="list-style-type: none"> Decreased food intake, glucose and HbA_{1c} concentrations Enhanced circulating and pancreatic insulin levels Improved glucose tolerance and glucose-induced insulin secretion Enhanced metabolic response to GIP and the glucose-lowering actions of insulin 	Hasib et al ⁷⁹

element have focussed on xenin-8 sequences, but utilisation of xenin-6 peptides, particularly xenin-6-psi, could offer distinct advantages over this approach. In terms of potential side effects of xenin-based therapeutics, the only notable reported side effect following xenin infusion in humans was mild diarrhoea.¹³ There is also lack of any obvious side effects in rodents following sustained xenin administration in numerous studies.^{54,76} Thus, xenin appears to have side-effect profile similar to that of established GLP-1 therapeutics, namely mild

gastrointestinal adverse events, with GLP-1 mimetics now well-established as important anti- obesity and -diabetes drugs in man.^{86,87} However, further dose-response studies are still required in human volunteers to uncover the complete adverse side effect profile of xenin. Ultimately, xenin-based therapies need to be further assessed in the human setting to confirm translatability of the many positive findings from pre-clinical trials,^{21,26,59,76,78,79,82,83} and progress benefits towards the clinic.

ORCID iD

Sarah L Craig  <https://orcid.org/0000-0002-3123-8834>

REFERENCES

- Feurle GE, Hamscher G, Kusiek R, Meyer HE, Metzger JW. Identification of xenin, a xenopsin-related peptide, in the human gastric mucosa and its effect on exocrine pancreatic secretion. *J Biol Chem*. 1992;267:22305-22309.
- Kageyama T, Ichinose M, Yonezawa S. Processing of the precursors to neurotensin and other bioactive peptides by cathepsin E. *J Biol Chem*. 1995;270:19135-19140.
- Hamscher G, Meyer HE, Feurle GE. Identification of proxenin as a precursor of the peptide xenin with sequence homology to yeast and mammalian coat protein α . *Peptides*. 1996;17:889-893.
- Feurle GE. Xenin – a review. *Peptides*. 1998;19:609-615.
- Chow VT, Quek HH. Alpha coat protein COPA (HEP-COP): presence of an Alu repeat in cDNA and identity of the amino terminus to xenin. *Ann Hum Genet*. 1997;61:369-373.
- Craig SL, Gault VA, Irwin N. Emerging therapeutic potential for xenin and related peptides in obesity and diabetes. *Diabetes Metab Res Rev*. 2018;34:e3006.
- Araki K, Tachibana S, Uchiyama M, Nakajima T, Yasuhara T. Isolation and structure of a new active peptide xenopsin on rat stomach strip and some biogenic amines in the skin of *Xenopus laevis*. *Chem Pharm Bull*. 1975;23:3132-3140.
- Hamscher G, Meyer HE, Metzger JW, Feurle GE. Distribution, formation, and molecular forms of the peptide xenin in various mammals. *Peptides*. 1995;16:791-797.
- Anlauf M, Weihe E, Hartschuh W, Hamscher G, Feurle GE. Localization of xenin-immunoreactive cells in the duodenal mucosa of humans and various mammals. *J Histochem Cytochem*. 2000;48:1617-1626.
- Khan D, Vasu S, Moffett RC, Gault VA, Flatt PR, Irwin N. Locally produced xenin and the neurotensinergic system in pancreatic islet function and β -cell survival. *Biol Chem*. 2017;399:79-92.
- Cline MA, Nandar W, Rogers JO. Xenin reduces feed intake by activating the ventromedial hypothalamus and influences gastrointestinal transit rate in chicks. *Behav Brain Res*. 2007;179:28-32.
- Kim ER, Mizuno TM. Xenin delays gastric emptying rate and activates the brainstem in mice. *Neurosci Lett*. 2010;481:59-63.
- Chowdhury S, Reeds DN, Crimmins DL, et al. Xenin-25 delays gastric emptying and reduces postprandial glucose levels in humans with and without type 2 diabetes. *Am J Physiol Gastrointest Liver Physiol*. 2014;306:G301-G309.
- Alexiou C, Zimmermann JP, Schick RR, Schusdziaara V. Xenin-a novel suppressor of food intake in rats. *Brain Res*. 1998;800:294-299.
- Leckstrom A, Kim ER, Wong D, Mizuno TM. Xenin, a gastrointestinal peptide, regulates feeding independent of the melanocortin signaling pathway. *Diabetes*. 2009;58:87-94.
- Martin CM, Parthasarathy V, Hasib A, et al. Biological activity and antidiabetic potential of C-terminal octapeptide fragments of the gut-derived hormone xenin. *PLoS One*. 2016;11:e0152818.
- Silvestre RA, Rodríguez-Gallardo J, Egido EM, Hernández R, Marco J. Stimulatory effect of xenin-8 on insulin and glucagon secretion in the perfused rat pancreas. *Regul Pept*. 2003;115:25-29.
- Taylor AI, Irwin N, McKillop AM, Patterson S, Flatt PR, Gault VA. Evaluation of the degradation and metabolic effects of the gut peptide xenin on insulin secretion, glycaemic control and satiety. *J Endocrinol*. 2010;207:87-93.
- Martin CM, Gault VA, McClean S, Flatt PR, Irwin N. Degradation, insulin secretion, glucose-lowering and GIP additive actions of a palmitate-derivatised analogue of xenin-25. *Biochem Pharmacol*. 2012;84:312-319.
- Martin CM, Parthasarathy V, Pathak V, Gault VA, Flatt PR, Irwin N. Characterisation of the biological activity of xenin-25 degradation fragment peptides. *J Endocrinol*. 2014;221:193-200.
- Gault VA, Martin CMA, Flatt PR, Parthasarathy V, Irwin N. Xenin-25[Lys(13) PAL]: a novel long-acting acylated analogue of xenin-25 with promising antidiabetic potential. *Acta Diabetol*. 2015;52:461-471.
- Parthasarathy V, Irwin N, Hasib A, et al. A novel chemically modified analogue of xenin-25 exhibits improved glucose-lowering and insulin-releasing properties. *Biochim Biophys Acta*. 2016;1860:757-764.
- Gobron B, Bouvard B, Vyavahare S, et al. Enteroendocrine K cells exert complementary effects to control bone quality and mass in mice. *J Bone Miner Res*. 2020;35:1363-1374.
- Wice BM, Reeds DN, Tran HD, et al. Xenin-25 amplifies GIP-mediated insulin secretion in humans with normal and impaired glucose tolerance but not type 2 diabetes. *Diabetes*. 2012;61:1793-1800.
- Wice BM, Wang S, Crimmins DL, et al. Xenin-25 potentiates glucose-dependent insulinotropic polypeptide action via a novel cholinergic relay mechanism. *J Biol Chem*. 2010;285:19842-19853.
- Craig SL, Gault VA, McClean S, Hamscher G, Irwin N. Effects of an enzymatically stable C-terminal hexapeptide fragment peptide of xenin-25, ψ -xenin-6, on pancreatic islet function and metabolism. *Mol Cell Endocrinol*. 2019;496:110523.
- Clemens A, Katsoulis S, Nustede R, et al. Relaxant effect of xenin on rat ileum is mediated by apamin-sensitive neurotensin-type receptors. *Am J Physiol*. 1997;272:G190-G196.
- Heuser M, Kleiman I, Pöpkén O, Nustede R, Post S. Evidence for non-neurotensin receptor-mediated effects of xenin (1-25)—focus on intestinal microcirculation. *Regul Pept*. 2002;107:23-27.
- Tanday N, Moffett RC, Gault VA, Flatt PR, Irwin N. Enzymatically stable analogue of the gut-derived peptide xenin on beta-cell transdifferentiation in high fat fed and insulin-deficient *Ins1^{Cre/+}; Rosa26-eYFP* mice. *Diabetes Metab Res Rev*. 2021;37:e3384.
- Feurle GE, Klein A, Hamscher G, Metzger JW, Schuurkes JA. Neurokinetic and myokinetic effects of the peptide xenin on the motility of the small and large intestine of Guinea pig. *J Pharmacol Exp Ther*. 1996;278:654-661.
- Cooke JH, Patterson M, Patel SR, et al. Peripheral and central administration of xenin and neurotensin suppress food intake in rodents. *Obesity*. 2009;17:1135-1143.
- Bhavya S, Lew PS, Mizuno TM. Central action of xenin affects the expression of lipid metabolism-related genes and proteins in mouse white adipose tissue. *Neuropeptides*. 2017;63:67-73.
- English A, Craig SL, Flatt PR, Irwin N. Individual and combined effects of GIP and xenin on differentiation, glucose uptake and lipolysis in 3T3-L1 adipocytes. *Biol Chem*. 2020;401:1293-1303.
- Knop FK, Vilsbøll T, Højberg PV, et al. Reduced incretin effect in type 2 diabetes: cause or consequence of the diabetic state? *Diabetes*. 2007;56:1951-1959.
- Nauck MA, Meier JJ. Incretin hormones: their role in health and disease. *Diabetes Obes Metab*. 2018;20 Suppl 1:5-21.
- Willard FS, Douros JD, Gabe MB, et al. Tirzepatide is an imbalanced and biased dual GIP and GLP-1 receptor agonist. *JCI Insight*. 2020;5:e140532.
- Mazella J, Béraud-Dufour S, Devader C, Massa F, Coppola T. Neurotensin and its receptors in the control of glucose homeostasis. *Front Endocrinol*. 2012;3:143.
- Kim ER, Lew PS, Spirkina A, Mizuno TM. Xenin-induced feeding suppression is not mediated through the activation of central extracellular signal-regulated kinase signaling in mice. *Behav Brain Res*. 2016;312:118-126.
- Kerbel B, Badal K, Sundararajan L, Blanco A, Unniappan S. Xenin is a novel anorexigen in goldfish (*Carassius auratus*). *PLoS One*. 2018;13:e0197817.
- Trümper A, Trümper K, Hörsch D. Mechanisms of mitogenic and anti-apoptotic signaling by glucose-dependent insulinotropic polypeptide in beta(INS-1)-cells. *J Endocrinol*. 2002;174:233-246.
- Trümper A, Trümper K, Trusheim H, Arnold R, Göke B, Hörsch D. Glucose-dependent Insulinotropic polypeptide is a growth factor for β (INS-1) cells by Pleiotropic signaling. *Mol Endocrinol*. 2001;15:1559-1570.
- Harada N, Inagaki N. Role of GIP receptor signaling in β -cell survival. *Diabetol Int*. 2017;8:137-138.
- Sarnobat D, Moffett RC, Gault VA, et al. Effects of long-acting GIP, xenin and oxyntomodulin peptide analogues on alpha-cell transdifferentiation in insulin-deficient diabetic *GluCreERT2; ROSA26-eYFP* mice. *Peptides*. 2020;125:170205.
- Rojas J, Chávez M, Olivar L, et al. Polycystic ovary syndrome, insulin resistance, and obesity: navigating the pathophysiological labyrinth. *Int J Reprod Med*. 2014;2014:719050.
- Temur M, Özün Özbay P, Aksun S, et al. Elevated circulating levels of xenopsin-related peptide-1 are associated with polycystic ovary syndrome. *Arch Gynecol Obstet*. 2017;296:841-846.
- Guclu YA, Sahin E, Aksit M. The relationship between elevated serum xenin and insulin resistance in women with polycystic ovary syndrome: a case-control study. *Gynecol Endocrinol*. 2019;35:960-964.
- Marques S, Pirogova E, Piva TJ. Evaluation of the use of therapeutic peptides for cancer treatment. *J Biomed Sci*. 2017;24:21.
- Fosgerau K, Hoffmann T. Peptide therapeutics: current status and future directions. *Drug Discov Today*. 2015;20:122-128.
- Di L. Strategic approaches to optimizing peptide ADME properties. *AAPS J*. 2015;17:134-143.
- Feurle GE, Meyer HE, Hamscher G. Metabolism and potency of xenin and of its reduced hexapeptide psi fragment in the dog. *Life Sci*. 2003;74:697-707.
- Werle M, Bernkop-Schnürch A. Strategies to improve plasma half life time of peptide and protein drugs. *Amino Acids*. 2006;30:351-367.
- Madsen K, Knudsen LB, Agersøe H, et al. Structure-activity and protraction relationship of long-acting glucagon-like peptide-1 derivatives: importance of fatty acid length, polarity, and bulkiness. *J Med Chem*. 2007;50:6126-6132.
- Goodwin D, Simerska P, Toth I. Peptides as therapeutics with enhanced bioactivity. *Curr Med Chem*. 2012;19:4451-4461.

54. Hasib A, Ng MT, Gault VA, et al. An enzymatically stable GIP/xenin hybrid peptide restores GIP sensitivity, enhances beta cell function and improves glucose homeostasis in high-fat-fed mice. *Diabetologia*. 2017;60:541-552.
55. Novakovic ZM, Leinung MC, Grasso P. [D-Leu-4]-OB3, an orally bioavailable leptin-related synthetic peptide insulin sensitizer: a study comparing the efficacies of [D-Leu-4]-OB3 and metformin on energy balance and glycemic regulation in insulin-deficient male Swiss Webster mice. *Peptides*. 2013;43:167-173.
56. Park JW, Kim SJ, Kwag DS, et al. Multifunctional delivery systems for advanced oral uptake of peptide/protein drugs. *Curr Pharm Des*. 2015;21:3097-3110.
57. Feurle GE, Heger M, Niebergall-Roth E, et al. Gastroenteropancreatic effects of xenin in the dog. *J Pept Res*. 1997;49:324-330.
58. Rodriguez-Gallardo J, Egido E, Gutierrez E, Garcí'a P, Silvestre RA, Marco J. Effect of xenin-8 on pancreatic hormone secretion in the rat pancreas. *Diabetologia*. 2001;44:A193.
59. Craig SL, Gault VA, Hamscher G, Irwin N. Ψ -xenin-6 enhances sitagliptin effectiveness, but does not improve glucose tolerance. *J Endocrinol*. 2020;245:219-230.
60. Irwin N, Flatt PR. New perspectives on exploitation of incretin peptides for the treatment of diabetes and related disorders. *World J Diabetes*. 2015;6:1285.
61. Coskun T, Sloop KW, Lohin C, et al. LY3298176, a novel dual GIP and GLP-1 receptor agonist for the treatment of type 2 diabetes mellitus: from discovery to clinical proof of concept. *Mol Metab*. 2018;18:3-14.
62. ClinicalTrials.gov. A study of tirzepatide (LY3298176) versus insulin degludec in participants with type 2 diabetes (SURPASS-3); 2020.
63. Pocai A. Action and therapeutic potential of oxyntomodulin. *Mol Metab*. 2013;3:241-251.
64. Irwin N, Pathak V, Flatt PR. A novel CCK-8/GLP-1 hybrid peptide exhibiting prominent insulinotropic, glucose-lowering, and satiety actions with significant therapeutic potential in high-fat-fed mice. *Diabetes*. 2015;64:2996-3009.
65. Mansur SA, Mieczkowska A, Flatt PR, et al. A new stable GIP-Oxyntomodulin hybrid peptide improved bone strength both at the organ and tissue levels in genetically-inherited type 2 diabetes mellitus. *Bone*. 2016;87:102-113.
66. Tschöp M, DiMarchi R. Single-molecule combinatorial therapeutics for treating obesity and diabetes. *Diabetes*. 2017;66:1766-1769.
67. Gault VA, Kerr BD, Harriott P, Flatt PR. Administration of an acylated GLP-1 and GIP preparation provides added beneficial glucose-lowering and insulinotropic actions over single incretins in mice with type 2 diabetes and obesity. *Clin Sci*. 2011;121:107-117.
68. Irwin N, McClean PL, Flatt PR. Comparison of the subchronic antidiabetic effects of DPP IV-resistant GIP and GLP-1 analogues in obese diabetic (ob/ob) mice. *J Pept Sci*. 2007;13:400-405.
69. Irwin N, McClean PL, Cassidy RS, et al. Comparison of the anti-diabetic effects of GIP- and GLP-1-receptor activation in obese diabetic (ob/ob) mice: studies with DPP IV resistant N-AcGIP and exendin(1-39)amide. *Diabetes Metab Res Rev*. 2007;23:572-579.
70. Parker JC, Irwin N, Lavery KS, et al. Metabolic effects of sub-chronic ablation of the incretin receptors by daily administration of (Pro3)GIP and exendin(9-39) amide in obese diabetic (ob/ob) mice. *Biol Chem*. 2007;388:221-226.
71. Gault VA, Bhat VK, Irwin N, Flatt PR. A novel Glucagon-like peptide-1 (GLP-1)/glucagon hybrid peptide with triple-acting agonist activity at glucose-dependent insulinotropic polypeptide, GLP-1, and glucagon receptors and therapeutic potential in high fat-fed mice. *J Biol Chem*. 2013;288:35581-35591.
72. Hornigold DC, Roth E, Howard V, et al. A GLP-1:CCK fusion peptide harnesses the synergistic effects on metabolism of CCK-1 and GLP-1 receptor agonism in mice. *Appetite*. 2018;127:334-340.
73. Bhat VK, Kerr BD, Flatt PR, Gault VA. A novel GIP-oxyntomodulin hybrid peptide acting through GIP, glucagon and GLP-1 receptors exhibits weight reducing and anti-diabetic properties. *Biochem Pharmacol*. 2013;85:1655-1662.
74. Pathak NM, Pathak V, Gault VA, McClean S, Irwin N, Flatt PR. Novel dual incretin agonist peptide with antidiabetic and neuroprotective potential. *Biochem Pharmacol*. 2018;155:264-274.
75. Tschöp MH, Finan B, Clemmensen C, et al. Unimolecular polypharmacy for treatment of diabetes and obesity. *Cell Metab*. 2016;24:51-62.
76. Craig SL, Perry RA, Vyavahare SS, et al. A GIP/xenin hybrid in combination with exendin-4 improves metabolic status in db/db diabetic mice and promotes enduring antidiabetic benefits in high fat fed mice. *Biochem Pharmacol*. 2020;171:113723.
77. Prasad RB, Groop L. Genetics of type 2 diabetes—pitfalls and possibilities. *Genes*. 2015;6:87-123.
78. Hasib A, Ng MT, Khan D, Gault VA, Flatt PR, Irwin N. A novel GLP-1/xenin hybrid peptide improves glucose homeostasis, circulating lipids and restores GIP sensitivity in high fat fed mice. *Peptides*. 2018;100:202-211.
79. Hasib A, Ng MT, Tanday N, et al. Exendin-4(Lys27 PAL)/gastrin/xenin-8-gln: a novel acylated GLP-1/gastrin/xenin hybrid peptide that improves metabolic status in obese-diabetic (ob/ob) mice. *Diabetes Metab Res Rev*. 2019;35:e3106.
80. Dalboge LS, Almholzt DLC, Neerup TSR, Vrang N, Jelsing J, Fosgerau K. The novel GLP-1-gastrin dual agonist ZP3022 improves glucose homeostasis and increases β -cell mass without affecting islet number in db/db mice. *J Pharmacol Exp Ther*. 2014;350:353-360.
81. Hasib A, Ng MT, Khan D, Gault VA, Flatt PR, Irwin N. Characterisation and antidiabetic utility of a novel hybrid peptide, exendin-4/gastrin/xenin-8-Gln. *Eur J Pharmacol*. 2018;834:126-135.
82. Craig SL, Gault VA, Flatt PR, Irwin N. The methionine aminopeptidase 2 inhibitor, TNP-470, enhances the antidiabetic properties of sitagliptin in mice by upregulating xenin. *Biochem Pharmacol*. 2021;183:114355.
83. Joharapurkar AA, Dhanesha NA, Jain MR. Inhibition of the methionine aminopeptidase 2 enzyme for the treatment of obesity. *Diabetes Metab Syndr Obes*. 2014;7:73-84.
84. Bräkenhielm E, Cao R, Gao B, et al. Angiogenesis inhibitor, TNP-470, prevents diet-induced and genetic obesity in mice. *Circ Res*. 2004;94:1579-1588.
85. White HM, Acton AJ, Considine RV. The angiogenic inhibitor TNP-470 decreases caloric intake and weight gain in high-fat fed mice. *Obesity (Silver Spring)*. 2012;20:2003-2009.
86. DeFronzo RA, Ratner RE, Han J, Kim DD, Fineman MS, Baron AD. Effects of exenatide (exendin-4) on glycemic control and weight over 30 weeks in metformin-treated patients with type 2 diabetes. *Diabetes Care*. 2005;28:1092-1100.
87. Nauck MA, Quast DR, Wefers J, Meier JJ. GLP-1 receptor agonists in the treatment of type 2 diabetes – state-of-the-art. *Mol Metab*. 2021;46:101102.